

LOCAL AND MICROLOCAL BMT REGULARITY VIA A CLASS OF FBI TRANSFORMS

RENAN DANTAS MEDRADO

ABSTRACT. In this talk I will present a characterization of local and microlocal BMT regularity (in the sense of Braun, Meise and Taylor, see [1]) for ultradistributions using a class of FBI transforms defined by weight functions (this class involves the weight functions used to define BMT regularity and includes the subclass of Sjostrand's FBI transforms used in [2]). Also, I will present a characterization of Iterates of Partial Differential Operators in the sense of BMT classes (called BMT vectors) and, as an application, I will present a relation between BMT regularity and BMT vectors.

Joint work with G. Hoepfner. Partially supported by FAPEAL.

REFERENCES

- [1] Braun, R. W.; Meise, R. and Taylor, B. A. Ultradifferentiable functions and Fourier analysis. *Res. Math.*, 17, 207–237, 1990.
- [2] Christ, M. Intermediate optimal Gevrey exponents occur. *Comm. Partial Differential Equations.*, 22, n. 3-4, 359–379, 1997.
- [3] Komatsu, H. A characterization of Real Analytic Functions. *Proc. Japan Acad.*, 1960.
- [4] Nelson, E. Analytic vectors. *Ann. of Math.*, 70, 572–615, 1959.

UNIVERSIDADE FEDERAL DE ALAGOAS (UFAL)

Email address: `renan.medrado@im.ufal.br`